

# Deconfinement phase transition and the quark condensate

Christian S. Fischer


TU Darmstadt / GSI

9th June 2009

C.F., arXiv:0904.2700 [hep-ph]

C.F. and J. Mueller, in preparation

# QCD phase transitions



- Chiral limit ( $M_{weak} = 0$ ): order parameter chiral condensate
- Heavy quarks ( $M_{weak} = \infty$ ): order parameter Polyakov-loop

# Lattice QCD vs. DSE/FRG: Complementary!

- Lattice simulations
  - ▶ Ab initio
  - ▶ Gauge invariant
- Functional approaches:
  - Dyson-Schwinger equations (DSE)
  - Functional renormalisation group (FRG)
    - ▶ Analytic solutions at small momenta
    - ▶ Space-Time-Continuum
    - ▶ Chiral symmetry: light quarks and mesons
    - ▶ Chemical potential: no sign problem

# QCD in covariant gauge

$$\mathcal{Z}_{\text{QCD}} = \int \mathcal{D}[\Psi, A, c] \exp \left\{ - \int_0^{1/T} dt \int d^3x \left( \bar{\Psi} (i\not{D} - m) \Psi - \frac{1}{4} (F_{\mu\nu}^a)^2 + \frac{(\partial A)^2}{2\xi} + \bar{c}(-\partial D)c \right) \right\}$$

Landau gauge ( $\xi = 0$ ) propagators in momentum space,  $q = (\vec{q}, \omega_q)$ :



$$D_{\mu\nu}^{\text{Gluon}}(q) = \frac{Z_T(q)}{q^2} P_{\mu\nu}^T(q) + \frac{Z_L(q)}{q^2} P_{\mu\nu}^L(q)$$



$$S^{\text{Quark}}(q) = \frac{1}{-i \vec{\gamma} \vec{q} A(q) - i \gamma_4 \omega_n C(q) + B(q)}$$

The Goal:

Gauge invariant information from gauge fixed functional approach

# The ordinary chiral condensate

The diagram shows an equation between three terms. On the left is a horizontal line with a shaded circle in the middle. Above it is a superscript  $-1$ . This is followed by an equals sign. To the right of the equals sign is a horizontal line with a superscript  $-1$  above it, followed by a plus sign. To the right of the plus sign is a loop diagram. The loop consists of a shaded circle at the top and a white circle at the bottom, connected by two wavy lines. The loop is attached to a horizontal line on the left and right.

- Consider DSE on torus with  $V = 1/T \times L^3$ 
  - spatial directions: **periodic** boundary conditions
  - temporal direction: **antiperiodic** boundary condition

C.F., arXiv:0904.2700 [hep-ph]

- Consider DSE in infinite volume/continuum

C.F. and J.Mueller, in preparation

- Order parameter for **chiral transition**:

$$\langle \bar{\psi}\psi \rangle = Z_2 N_c \frac{T}{L^3} \text{Tr}_D \sum_{\vec{p}, \omega_p} S(p_{\vec{p}, \omega_p})$$


# The dual condensate I

Consider general  $U(1)$ -valued boundary conditions in temporal direction for quark fields  $\psi$ :

$$\psi(\vec{x}, 1/T) = e^{i\varphi} \psi(\vec{x}, 0)$$

Matsubara frequencies:  $\omega_p(n_t) = (2\pi T)(n_t + \varphi/2\pi)$

Lattice:




E. Bilgici, F. Bruckmann, C. Gattringer and C. Hagen, PRD **77** (2008) 094007..

# The dual condensate II

Relation of condensate to loops of link variables  $U_\mu(x)$ :

$$\langle \bar{\psi} \psi \rangle_\varphi = \text{Tr} [m + D_\varphi]^{-1} = \frac{1}{Vm} \sum_{l \in \mathcal{L}} \frac{e^{i\varphi n(l)}}{(2am)^{|l|}} \text{Tr}_c \prod_{(x,\mu) \in l} s(l) U_\mu(x).$$

- geometric series of inverse staggered Dirac operator
- winding number  $n(l)$  of loop  $l$  around temporal direction



# The dual condensate III

Then define dual condensate  $\Sigma_n$ :

$$\Sigma_n = - \int_0^{2\pi} \frac{d\varphi}{2\pi} e^{-i\varphi n} \langle \bar{\psi}\psi \rangle_\varphi$$

- $n = 1$  projects out loops with  $n(l) = 1$ : dressed Polyakov loop
- transforms under center transformation exactly like ordinary Polyakov loop
- $\Sigma_1$  is order parameter for center symmetry/deconfinement

E. Bilgici, F. Bruckmann, C. Gattringer and C. Hagen, PRD **77** (2008) 094007.

- $\Sigma_1$  is accessible with functional methods

C.F., arXiv:0904.2700 [hep-ph]


# Input into quark-DSE

$$\text{Diagram with shaded circle}^{-1} = \text{Diagram with plain line}^{-1} + \text{Diagram with loop}^{-1}$$


- $T$ -dependent gluon propagator from lattice data

Cucchieri, Maas, Mendes, PRD75 (2007)

transverse gluon dressing function





longitudinal gluon dressing function




- $T$ -dependent ansatz for quark-gluon vertex

# Lattice results



E. Bilgici, F. Bruckmann, C. Gattringer and C. Hagen, PRD **77** (2008) 094007..


# Results: angular dependence of condensate



$$\Delta(\varphi) \equiv \langle \bar{\psi}\psi \rangle_{\varphi} = \text{Tr} [m + D_{\varphi}]^{-1} = \frac{1}{Vm} \sum_{l \in \mathcal{L}} \frac{e^{i\varphi n(l)}}{(2am)^{|l|}} \text{Tr}_c \prod_{(x,\mu) \in l} s(l) U_{\mu}(x).$$

- Smaller mass: more contributions from loop with larger  $q(l)$ !


# Results: angular dependence in chiral limit



- Chiral limit: need continuum DSEs
- Chiral limit: all terms of expansion contribute
- Width of plateau is  $T$ -dependent!

C.F. and Jens Mueller, in preparation

# Results: dressed Polyakov loop $\Sigma_1$



- Deconfinement transition from functional methods


C.F., arXiv:0904.2700 [hep-ph]

C.F. and Jens Mueller, in preparation

- other method: Polyakov loop potential from ghost and glue

J. Braun, H. Gies and J. M. Pawłowski, arXiv:0708.2413 [hep-th].

# Results: dual skalar quark dressing $\Sigma_B$





Definition of dual skalar quark dressing  $\Sigma_B$ :

$$\Sigma_B = - \int_0^{2\pi} \frac{d\varphi}{2\pi} e^{-i\varphi n} B(0)$$

F. Synatschke, A. Wipf and K. Langfeld, PRD **77** (2008) 114018

C.F. and Jens Mueller, in preparation

# Results: chiral limit



- quark mass  $m(20\text{GeV}^2) = 10$  MeV vs. chiral limit
- **chiral limit: transition temperatures coincide**

C.F. and Jens Mueller, in preparation

## Techniques:

- Dual condensate → deconfinement order parameter
- Dual skalar quark dressing → deconfinement order parameter
- Calculable with functional methods!

## Results for $D_\chi$ SB and Deconfinement:

- Different transition temperatures at finite quark mass
- Same transition temperatures in chiral limit

# This work was supported by:

## Helmholtz Young Investigator Group "Nonperturbative Phenomena in QCD"



TECHNISCHE  
UNIVERSITÄT  
DARMSTADT



HELMHOLTZ  
| GEMEINSCHAFT



 **LOEWE** – Landes-Offensive zur Entwicklung  
Wissenschaftlich-ökonomischer Exzellenz



Helmholtz-Alliance: Extremes of density and temperature; cosmic matter in the laboratory

# Gluon and Quark-Gluon-Vertex

Fit function for gluon:

$$Z_{T,L}(\vec{q}, \omega_q, T) = \frac{q^2 \Lambda^2}{(q^2 + \Lambda^2)^2} \left\{ \left( \frac{c}{q^2 + \Lambda^2 a_{T,L}(T)} \right)^2 + \frac{q^2}{\Lambda^2} \left( \frac{\beta_0 \alpha(\mu) \ln[q^2/\Lambda^2 + 1]}{4\pi} \right)^\gamma \right\}$$

Ansatz for Quark-Gluon-Vertex:

$$\Gamma_\nu(q, k, p) = \tilde{Z}_3 \left( \delta_{4\nu} \gamma_4 \frac{C(k) + C(p)}{2} + \delta_{j\nu} \gamma_j \frac{A(k) + A(p)}{2} \right) \times \left( \frac{d_1}{d_2 + q^2} + \frac{q^2}{\Lambda^2 + q^2} \left( \frac{\beta_0 \alpha(\mu) \ln[q^2/\Lambda^2 + 1]}{4\pi} \right)^{2\delta} \right).$$