# Charge Fluctuations and Correlations at zero and non zero Density

Christian Schmidt
for the RBC-Bielefeld Collaboration

Lattice 2007
REGENSBURG

# Outline

- Introduction
- Taylor expansion in B, S, Q chemical potentials

  $\longrightarrow$ generalized susceptibilities at $\mu_{B,S,Q} = 0$
- Baryon, strangeness and charge fluctuations at $\mu_B > 0 \quad (\mu_S = \mu_Q = 0)$
- Dens matter with constraints on charges

  $\longrightarrow$ the HIC fireball $(\langle n_S \rangle = 0)$

  $\longrightarrow$ flavor symmetric matter $(\langle n_q \rangle = \langle n_s \rangle)$
- Correlations between charges at $\mu_B > 0$

# Introduction (1)

$$\mu_S = \mu_Q = 0$$

- Fluctuations of B, S, Q can be measured experimentally
- LGT at $\mu = 0$
  $\longrightarrow$ RHICH, LHC
- LGT at $\mu > 0$
  $\longrightarrow$ RHICH at low energies, FAIR@GSI




$\sim$ few times nuclear matter density

# Introduction (2)

At $\mu = 0$ and $m_{u,d}$ small, we expect O(4)-critical behavior

- scaling field:

$$t = \left|\frac{T - T_c}{T_c}\right| + A\left(\left(\frac{\mu_u}{T_c}\right)^2 + \left(\frac{\mu_d}{T_c}\right)^2\right) + B\frac{\mu_u}{T_c}\frac{\mu_d}{T_c}$$

- singular part:

$$f_s(T, \mu_u, \mu_d) = b^{-1} f_s(t b^{1/(2-\alpha)}) \sim t^{(2-\alpha)}$$

- O(4)/O(2): $\alpha < 0$, small

$\left\langle (\delta n_{u,d})^2 \right\rangle \longrightarrow$ dominated by T dependence of regular part

$\left\langle (\delta n_{u,d})^4 \right\rangle \longrightarrow$ develops a cusp!






# Taylor expansion in $\mu_{B,S,Q}$

QCD is naturally formulated with quark chemical potentials $\mu_{u,d,s}$

- we set $\mu_u \equiv \mu_d$ and start from Taylor expansion of the pressure

$$\frac{p}{T^4} = \frac{1}{VT^3} \ln Z(V,T,\mu_q,\mu_s) = \sum_{i,j} c_{i,j}^{q,s} \left(\frac{\mu_q}{T}\right)^i \left(\frac{\mu_s}{T}\right)^j$$

- use unbiased, noisy estimators to calculate $c_{i,j}^{q,s}$
  $\longrightarrow$ see Talk by C. Miao

- Line of constant physics: $m_q = m_s/10$
  (physical strange quark mass)

- measure currently up to $\mathcal{O}(\mu^8) \longleftrightarrow (N_t = 4)$
  $\mathcal{O}(\mu^4) \longleftrightarrow (N_t = 6)$

# Taylor expansion in $\mu_{B,S,Q}$

QCD is naturally formulated with quark chemical potentials $\mu_{u,d,s}$

- we set $\mu_u \equiv \mu_d$ and start from Taylor expansion of the pressure

$$\frac{p}{T^4} = \frac{1}{VT^3} \ln Z(V,T,\mu_q,\mu_s) = \sum_{i,j} c_{i,j}^{q,s} \left(\frac{\mu_q}{T}\right)^i \left(\frac{\mu_s}{T}\right)^j$$

- expansion coefficients $c_{i,j}^{q,s}$ are related to B,S,Q-fluctuations

$$\begin{aligned} n_B &= \frac{\partial(p/T^4)}{\partial(\mu_B/T)} = \frac{1}{3}(n_u + n_d + n_s) \\ n_S &= \frac{\partial(p/T^4)}{\partial(\mu_S/T)} = -n_s \\ n_Q &= \frac{\partial(p/T^4)}{\partial(\mu_Q/T)} = \frac{1}{3}(2n_u - n_d - n_s) \end{aligned}$$

$$\begin{aligned} \mu_u &= \frac{1}{3}\mu_B + \frac{2}{3}\mu_Q \\ \mu_d &= \frac{1}{3}\mu_B - \frac{1}{3}\mu_Q \\ \mu_s &= \frac{1}{3}\mu_B - \frac{1}{3}\mu_Q - \mu_S \end{aligned}$$

- choice of $\mu_u \equiv \mu_d$ is equivalent to $\mu_Q \equiv 0$

# Hadronic fluctuations $(\mu_B = 0)$

In general we have:

- $\dfrac{\partial^p (p/T^4)}{\partial(\mu_X/T)^p}$ related to $< n_X^p >_{\mu=0}$
- $\dfrac{\partial^{p+q} (p/T^4)}{\partial(\mu_X/T)\partial(\mu_Y/T)^q}$ related to $< n_X^p n_Y^q >_{\mu=0}$

to be more precise:

$$
\begin{aligned}
2c_2^X = \frac{\partial^2 (p/T^4)}{\partial(\mu_X/T)^2} &= \frac{1}{VT^3}\left\langle (\delta N_X)^2 \right\rangle = \frac{1}{VT}\left\langle N_X^2 \right\rangle \\
24c_4^X = \frac{\partial^4 (p/T^4)}{\partial(\mu_X/T)^4} &= \frac{1}{VT^3}\left(\left\langle (\delta N_X)^4 \right\rangle - 3\left\langle (\delta N_X^2) \right\rangle^2\right)_{\mu=0} \\
&= \frac{1}{VT^3}\left(\left\langle N_X^4 \right\rangle - 3\left\langle N_X^2 \right\rangle^2\right)_{\mu=0} \\
c_{11}^{XY} = \frac{\partial^2 (p/T^4)}{\partial(\mu_X/T)\partial(\mu_Y/T)} &= \frac{1}{VT^3}\left(\left\langle N_X N_Y \right\rangle - \left\langle N_X \right\rangle \left\langle N_Y \right\rangle\right)_{\mu=0}
\end{aligned}
$$

# Baryon number fluctuations $(\mu_B = 0)$







# Strangeness fluctuations $(\mu_B = 0)$






# Electric charge fluctuations $(\mu_B = 0)$










- Ratio is sensitive to charge quanta in the QGP

→ consistent with a gas of quasi-free quarks and gluons.

red: RBC-Bielefeld, 2+1 flavor, preliminary

blue: Ejiri, Karsch, Redlich, PLB 633 (2006) 275

# Baryon number and strangeness fluctuations at $\mu_B > 0 \quad (\mu_S = 0)$

$$\frac{\chi_B}{T^2} = 2c_2^B + 12c_4^B \left(\frac{\mu_B}{T}\right)^2$$




$$\frac{\chi_S}{T^2} = 2c_2^B + 2c_{22}^{BS} \left(\frac{\mu_B}{T}\right)^2$$




→ large baryon number fluctuations

→ enhanced strangeness fluctuations (~ factor 3 at Tc)

# The HIC case: $\langle N_s \rangle \equiv 0$

- Solving order by order:

$$
\begin{aligned}
N_s(\mu_q, \mu_q) &= \sum_{i,j} (j+1) c^{qs}_{i(j+1)} \mu_q^i \mu_s^j \\
&= c^{qs}_{11}\mu_q + 2c^{qs}_{02}\mu_s \\
&+ c^{qs}_{31}\mu_q^3 + 2c^{qs}_{22}\mu_q^2\mu_s + 3c^{qs}_{13}\mu_q\mu_s^2 + 4c^{qs}_{04}\mu_s^3 \\
&+ \mathcal{O}\left(\mu^5\right) \\
&\equiv 0
\end{aligned}
$$

- Solution:

$$
\mu_s(\mu_q) = \sum_i d_i \mu_q^i = \left(-\frac{c^{qs}_{11}}{2c^{qs}_{02}}\right)\mu_q + \left(\frac{2c^{qs}_{04}c^{qs\,3}_{11} - 3c^{qs}_{02}c^{qs\,2}_{11}c^{qs}_{13} + 4c^{qs\,2}_{02}c^{qs}_{11}c^{qs}_{22} - 4c^{qs\,3}_{02}c^{qs}_{31}}{8c^{qs\,4}_{02}}\right)\mu_q^3 + \mathcal{O}\left(\mu_q^5\right)
$$

$\longrightarrow$ new expansion Coefficients: $\hat{c}^q_i$

$$
\begin{aligned}
\frac{\Delta p}{T^4} &= \sum_i \hat{c}^q_i \mu_q^i \equiv \sum_{i,j} c_{ij}\mu^i \left(\sum_k d_k \mu_q^k\right)^j \\
&= \left(c^{qs}_{20} - \frac{c^{qs\,2}_{11}}{4c^{qs}_{02}}\right)\mu_q^2 + \left(c^{qs}_{40} + \frac{c^{qs}_{04}c^{qs\,4}_{11}}{16c^{qs\,4}_{02}} - \frac{c^{qs\,3}_{11}c^{qs}_{13}}{8c^{qs\,3}_{02}} + \frac{c^{qs\,2}_{11}c^{qs}_{22}}{4c^{qs\,2}_{02}} - \frac{c^{qs}_{11}c^{qs}_{31}}{2c^{qs}_{02}}\right)\mu_q^4 + \mathcal{O}(\mu_q^6)
\end{aligned}
$$

# The HIC case: $\langle N_s \rangle \equiv 0$

$\langle N_Q \rangle \equiv 0$ should be constrained as well

- Solving order by order:

$$
\begin{aligned}
N_s(\mu_q, \mu_q) &= \sum_{i,j} (j+1) c^{qs}_{i(j+1)} \mu_q^i \mu_s^j \\
&= c^{qs}_{11}\mu_q + 2c^{qs}_{02}\mu_s \\
&+ c^{qs}_{31}\mu_q^3 + 2c^{qs}_{22}\mu_q^2\mu_s + 3c^{qs}_{13}\mu_q\mu_s^2 + 4c^{qs}_{04}\mu_s^3 \\
&+ \mathcal{O}\left(\mu^5\right) \\
&\equiv 0
\end{aligned}
$$

- Solution:

$$
\mu_s(\mu_q) = \sum_i d_i \mu_q^i = \left(-\frac{c^{qs}_{11}}{2c^{qs}_{02}}\right)\mu_q + \left(\frac{2c^{qs}_{04}c^{qs\,3}_{11} - 3c^{qs}_{02}c^{qs\,2}_{11}c^{qs}_{13} + 4c^{qs\,2}_{02}c^{qs}_{11}c^{qs}_{22} - 4c^{qs\,3}_{02}c^{qs}_{31}}{8c^{qs\,4}_{02}}\right)\mu_q^3 + \mathcal{O}\left(\mu_q^5\right)
$$

$\longrightarrow$ new expansion Coefficients: $\hat{c}^q_i$

$$
\begin{aligned}
\frac{\Delta p}{T^4} &= \sum_i \hat{c}^q_i \mu_q^i \equiv \sum_{i,j} c_{ij}\mu^i \left(\sum_k d_k \mu_q^k\right)^j \\
&= \left(c^{qs}_{20} - \frac{c^{qs\,2}_{11}}{4c^{qs}_{02}}\right)\mu_q^2 + \left(c^{qs}_{40} + \frac{c^{qs}_{04}c^{qs\,4}_{11}}{16c^{qs\,4}_{02}} - \frac{c^{qs\,3}_{11}c^{qs}_{13}}{8c^{qs\,3}_{02}} + \frac{c^{qs\,2}_{11}c^{qs}_{22}}{4c^{qs\,2}_{02}} - \frac{c^{qs}_{11}c^{qs}_{31}}{2c^{qs}_{02}}\right)\mu_q^4 + \mathcal{O}(\mu_q^6)
\end{aligned}
$$

# The HIC case: $\langle N_s \rangle \equiv 0$




## baryon fluctuations

$$\frac{\hat{\chi}_B}{T^2} = 2\hat{c}_2^B + 12\hat{c}_4^B \left(\frac{\mu_B}{T}\right)^2$$



# The HIC case: $\langle N_s \rangle \equiv 0$

## baryon fluctuations

$$\frac{\hat{\chi}_B}{T^2} = 2\hat{c}_2^B + 12\hat{c}_4^B \left(\frac{\mu_B}{T}\right)^2$$










$$\frac{\hat{\chi}_B(\mu_B,T)}{\hat{\chi}_B^{sb}(\mu_B)} \Big/ \frac{\chi_B(\mu_B,T)}{\chi_B^{sb}(\mu_B)}$$

→ modifications below Tc ~ 30%

# The HIC case: $\langle N_s \rangle \equiv 0$




## strangeness fluctuations

$$\frac{\hat{\chi}_S}{T^2} = 2\hat{c}_2^B + 2\hat{c}_{22}^{BS}\left(\frac{\mu_B}{T}\right)^2$$



# The HIC case: $\langle N_s \rangle \equiv 0$




## strangeness fluctuations

$$\frac{\hat{\chi}_S}{T^2} = 2\hat{c}_2^B + 2\hat{c}_{22}^{BS}\left(\frac{\mu_B}{T}\right)^2$$







$$\frac{\hat{\chi}_S(\mu_B,T)}{\hat{\chi}_S^{sb}(\mu_B)} \Big/ \frac{\chi_S(\mu_B,T)}{\chi_S^{sb}(\mu_B)}$$

→ deviations up to 25% below Tc

# The flavor symmetric case:

- Solving order by order:

$$\langle N_s(\mu_q,\mu_s)\rangle \equiv \frac{1}{2}\langle N_q(\mu_q,\mu_s)\rangle$$

- Solution:

$$\begin{aligned}\mu_s(\mu_q) &= \sum_i d_i \mu_q^i = \left(\frac{c_{20}-c_{11}}{2c_{02}-c_{11}/2}\right)\mu_q \\ &+\left(\frac{64c_{04}(c_{11}-c_{20})^3}{(-4c_{02}+c_{11})^4}+\frac{24c_{13}(c_{11}-c_{20})^2}{(-4c_{02}+c_{11})^3}+\frac{8c_{22}(c_{11}-c_{20})}{(-4c_{02}+c_{11})^2}+\frac{2c_{31}}{(-4c_{02}+c_{11})}\right)\mu_q^3 \\ &+\mathcal{O}(\mu_q^5)\end{aligned}$$




→ Eventually of interest for cosmology and neutron / quark stars

# Correlations of S,B and S,Q




full: RBC-Bielefeld, nf=2+1, preliminary

open: Gavai, Gupta, nf=2, partially quenched

- Small quenching effect

$\longrightarrow$ Normalization effect ?

$$\frac{\langle N_S N_X \rangle}{\langle N_S^2 \rangle} = \frac{f_X c_{11}^{qs} + 2c_s^s}{2c_2^s}$$

$$\frac{1}{T}\left(\langle N_B N_S \rangle - \langle N_B \rangle \langle N_S \rangle\right) = \hat{c}_{11}^{BS} + 3\hat{c}_{31}^{BS}\left(\frac{\mu_B}{T}\right)^2$$




$\longrightarrow$ Correlations increase for $\mu_B > 0$

# Conclusions

- Taylor expansion method at 2+1 flavor have rich HIC phenomenology.
- All results on fluctuations and correlations develop a peak with increasing chemical potential and are consistent with a gas of quasi-free quarks, already at 1.2-1.5 Tc.
- The strangeness can be easily constrained to zero, which is more relevant for HIC.
- In future: the electric charge density should be constrained to zero as well (for the HIC case).
- The density of all quarks can be constrained to be equal (interesting for cosmology, stars?)